



TITLE:

Asymptotic expansions of the non-holomorphic Eisenstein series (Diophantine Problems and Analytic Number Theory)

AUTHOR(S):

Noda, Takumi

CITATION:

Noda, Takumi. Asymptotic expansions of the non-holomorphic Eisenstein series (Diophantine Problems and Analytic Number Theory). 数理解析研究所講究録 2003, 1319: 29-32

ISSUE DATE:

2003-05

URL:

<http://hdl.handle.net/2433/43046>

RIGHT:

Asymptotic expansions of the non-holomorphic Eisenstein series

Takumi Noda 野田 工

College of Engineering Nihon University 日本大学 工学部

Abstract

In this report we describe one asymptotic expansion of the non-holomorphic Eisenstein series using Airy functions. We might expect that the property of the Eisenstein series on the complex parameter is simple because of the "good behavior" of its constant terms. But the fact is that it is not so, and this makes our study interesting. We will find an analogy between Eisenstein series and the square of the Riemann zeta-function in the point of view of the asymptotic expansion.

1 Eisenstein series

Let $k \geq 0$ be an even integer and H be the upper half plane. The non-holomorphic Eisenstein series for $SL_2(\mathbb{Z})$ is defined by

$$E_k(z, s) = y^s \sum_{\{c, d\}} (cz + d)^{-k} |cz + d|^{-2s}. \quad (1)$$

Here $z = x + \sqrt{-1}y \in H$, $s \in \mathbb{C}$ and the summation is taken over $\begin{pmatrix} * & * \\ c & d \end{pmatrix}$, a complete system of representation of $\{ \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \in SL_2(\mathbb{Z}) \} \setminus SL_2(\mathbb{Z})$. The right-hand side of (1) converges absolutely and locally uniformly on $\{(z, s) \mid z \in H, \Re(s) > 1 - \frac{k}{2}\}$, and $E_k(z, s)$ has a meromorphic continuation to the whole s -plane.

In this report we consider the Eisenstein series $E(z, s) = E_0(z, s)$. Let $i = \sqrt{-1}$, $s = \sigma + it \in \mathbb{C}$ and $\zeta(s)$ be the Riemann zeta-function. For $\Re(s) > 1$, $\zeta(2s)E(z, s)$ is expressed by

$$\zeta(2s)E(z, s) = \frac{1}{2} \sum_{\substack{(c, d) \in \mathbb{Z}^2 \\ (c, d) \neq (0, 0)}} \frac{y^s}{|cz + d|^{2s}}.$$

The Fourier expansion is as follows;

$$\begin{aligned} \zeta(2s)E(z, s) &= \zeta(2s)y^s + \sqrt{\pi}\zeta(2s-1)\frac{\Gamma(s-\frac{1}{2})}{\Gamma(s)}y^{1-s} \\ &\quad + 4\frac{\pi^s}{\Gamma(s)}\sqrt{y} \sum_{n=1}^{\infty} n^{s-\frac{1}{2}} \sigma_{1-2s}(n) K_{s-\frac{1}{2}}(2\pi ny) \cos(2\pi nx), \end{aligned} \quad (2)$$

$$\sigma_s(l) = \sum_{d|l, d>0} d^s,$$

and $K_\nu(\tau)$ ($\nu, \tau \in \mathbb{C}$) is the modified Bessel function defined by the integral

$$K_\nu(\tau) = \frac{1}{2} \int_0^\infty u^{\nu-1} \exp\left(-\frac{1}{2}\tau\left(u + \frac{1}{u}\right)\right) du.$$

It is well-known that the Fourier expansion (2) gives the holomorphic continuation of $\zeta(2s)E(z, s)$ to the whole s -plane except for the simple pole at $s = 1$, and gives the functional equation

$$\pi^{-s}\Gamma(s)\zeta(2s)E(z, s) = \pi^{-1+s}\Gamma(1-s)\zeta(2-2s)E(z, 1-s).$$

2 Asymptotic expansion

Based on Olver's works [4]-[7], Balogh ([1], [2]) gave one uniform asymptotic expansion of K_{it} ($t \in \mathbb{R}$) for large values t using Airy functions. The Airy function is defined by

$$\text{Ai}(\xi) = \frac{1}{\pi} \int_0^\infty \cos\left(\frac{1}{3}u^3 + \xi u\right) du = \frac{1}{\sqrt{3}\pi} \xi^{\frac{1}{2}} K_{\frac{1}{3}}\left(\frac{2}{3}\xi^{\frac{3}{2}}\right).$$

The uniform asymptotic expansion due to Balogh is as follows;

$$K_{it}(\tau) = \frac{\sqrt{2}\pi}{t^{\frac{1}{3}}} \exp\left(-\frac{\pi}{2}t\right) \left(\frac{\rho}{r^2-1}\right)^{\frac{1}{4}} \left\{ \text{Ai}(\xi) \left(1 + \sum_{k=1}^m t^{-2k} A_k(\rho)\right) + t^{-\frac{4}{3}} \text{Ai}'(\xi) \sum_{k=0}^{m-1} t^{-2k} B_k(\rho) + \varepsilon_{2m+1} \right\}, \quad (3)$$

where $t \in \mathbb{R}$, $\tau \in \mathbb{C}$, and r, ρ, ξ are parameters such that

$$r = \tau/t, \quad \frac{2}{3}\rho^{\frac{3}{2}} = (r^2 - 1)^{\frac{1}{2}} - \text{arcsec } r, \quad \xi = t^{\frac{2}{3}}\rho.$$

The coefficients are defined in [4] and [5] by

$$\begin{cases} A_k(\rho) = \sum_{l=0}^{2k} (-1) b_l \rho^{-\frac{3}{2}l} U_{2k-l} \\ \rho^{\frac{1}{2}} B_k(\rho) = \sum_{l=0}^{2k+1} (-1) a_l \rho^{-\frac{3}{2}l} U_{2k-l+1}. \end{cases}$$

Here a_l, b_l are real coefficients and U_k is a polynomial in $\frac{1}{\sqrt{r^2-1}}$. The error term ε_{2m+1} is defined in [6] and [7].

The asymptotic expansion (3) is useful near the transition point $t = \tau$. Expansions for the cases $\tau/t \not\sim 1$ or $\tau - t = o(\tau^{\frac{1}{3}})$ are already obtained by using saddle point method. (See [8].)

In the following we assume $\sigma = \frac{1}{2}$. Applying asymptotic expansions of the Bessel function to the Eisenstein series, we have the following theorem.

Theorem 1 *For any positive constant $\varepsilon > 0$, we have*

$$\begin{aligned}
 & \zeta(1+2it)E(z, \tfrac{1}{2}+it) \\
 &= 4\pi^{it}\sqrt{y} \sum_{n=1}^{N_1} n^{-it} \sigma_{2it}(n) \{t^2 - (2\pi ny)^2\}^{-\frac{1}{4}} \cos(2\pi nx) \left\{ \sqrt{2} \sin(f(t, n)) + O\left(\frac{1}{2\pi ny}\right) \right\} \\
 & \quad + 4\pi^{1+it}\sqrt{y} \sum_{n=N_1+1}^{N_2} n^{-it} \sigma_{2it}(n) \{t^2 - (2\pi ny)^2\}^{-\frac{1}{4}} |\xi|^{\frac{1}{4}} \cos(2\pi nx) \\
 & \quad \times \left\{ \text{Ai}(\xi) \left(1 + t^{-2} A_1(\rho) - \xi^{\frac{1}{2}} t^{-\frac{4}{3}} B_0(\rho)\right) + \exp(-\tfrac{2}{3}\xi^{\frac{3}{2}}) (1 + |\xi|)^{-\frac{1}{4}} O\left(\frac{1}{t}\right) \right\} \\
 & \quad \times \left\{ 1 + O\left(\frac{1}{t}\right) \right\} + O(t^\varepsilon).
 \end{aligned} \tag{4}$$

Here

$$N_1 = \left\lfloor \frac{1}{1+\delta} \cdot \frac{t}{2\pi y} \right\rfloor, \quad N_2 = \left\lfloor \frac{t}{2y} \right\rfloor$$

for some positive constant $\delta > 0$ and

$$f(t, n) = t \operatorname{arccosh} \left(\frac{t}{2\pi ny} \right) - \{t^2 - (2\pi ny)^2\}^{\frac{1}{2}} + \frac{1}{4}\pi.$$

In (4), we are able to describe error terms more precisely. In that sense we call (4) the asymptotic expansion of the Eisenstein series.

3 Remark

Let $D < 0$ be the discriminant of an imaginary quadratic field K and $\zeta_K(s)$ be the Dedekind zeta-function of K . Let $h(D)$ be the ideal class number of K and $f_1, f_2, \dots, f_{h(D)}$ be the equivalence classes of binary quadratic forms of discriminant D . Then we have

$$\sum_{i=1}^{h(D)} \zeta(2s) E(z(f_i), s) = 2^{-s-1} w |D|^{s/2} \zeta_K(s).$$

Here $z(f) = (-b + \sqrt{D})/2a \in H$ is the associated root of the quadratic form $f(X, Y) = aX^2 + bXY + cY^2$, and w is the number of roots of unity in K . (Cf. [9], [10, Sect.8, Sect.11].) It is also well known that $\zeta_K(s) = \zeta(s)L(s)$ for the L -function with Kronecker's symbol. This shows that Theorem 1 is one approach to the study of the product of classical zeta-functions. Especially we will find an analogy between Theorem 1 and formulas of Voronoi-Atkinson type for $\zeta^2(s)$ proved by Jutila [3].

References

- [1] C. B. Balogh, *Uniform asymptotic expansions of the modified Bessel function of the third kind of large imaginary order*, Bull.Amer. Math. Soc., 72, 1966, p40-43.
- [2] C. B. Balogh, *Asymptotic expansions of the modified Bessel function of the third kind of imaginary order*, SIAM-J.-Appl.-Math. 15, 1967, p1315-1323.
- [3] M. Jutila, *Transformation formulae for Dirichlet polynomials*, J. Number Theory 18, 1984, p135-156.
- [4] F. W. J. Olver, *The asymptotic solutions of linear differential equations of the second order for large values of a parameter*, Philos. Trans. Roy. Soc. London, Ser. A, 247, 1954, p307-327.
- [5] F. W. J. Olver, *The asymptotic expansions of Bessel functions of large order*, Ibid., 247, 1954, p328-368.
- [6] F. W. J. Olver, *Error bounds for first approximations in turning-point problems* SIAM-J.-Appl.-Math. 11, 1963, p748-772.
- [7] F. W. J. Olver, *Error bounds for asymptotic expansions in turning-point problems*, Ibid., 12, 1964, p200-214.
- [8] G. N. Watson, *A Treatise on the theory of Bessel functions (2nd edition)*, Cambridge University Press, 1944.
- [9] D. Zagier, *Eisenstein series and the Riemann zeta-function* (in *Automorphic forms, Representation theory and Arithmetic*: edited by S. Gelbert), Springer, 1981. p275-301.
- [10] D. Zagier, *Zetafunktionen und quadratische Körper*, Springer, 1981.